


Claims

1. Compound of the general formula I

5



in which


10

R^{11} denotes H, F, Cl, Br, I, CN, aryl, heterocyclyl or a halogenated or unsubstituted alkyl radical having 1 to 15 carbon atoms, where, in addition, one or more CH_2 groups in this radical may each be replaced, independently of one another, by $-C\equiv C-$, $-CH=CH-$, $-O-$, $-CO-$, $-CO-O-$ or $-O-CO-$ in such a way that O atoms are not linked directly to one another;

15

A

stands for



20

a

is 0, 1 or 2;

25

 Z^{11}


represents a single bond, $-CH_2-CH_2-$, $-CF_2-CF_2-$, $-CF_2-CH_2-$, $-CH_2-CF_2-$, $-CH_2-O-$, $-O-CH_2-$, $-CF_2-O-$ or $-O-CF_2-$;

W


denotes $>CH-$ or $>C=$;

30

B and D, independently of one another, stand for



5



b and d, independently of one another, are 0 or 1;

10

Y^{11} denotes $=O$, $=C(SR^{12})(SR^{13})$, $=CF_2$, $-H$, $-F$, $-Cl$, $-Br$, $-I$, $-CN$, $-OH$, $-SH$, $-CO-R^{14}$, $-OSO_2R^{15}$, $-C(=S^+R^{12})(-SR^{13})X^-$, $-B(OR^{16})(OR^{17})$, $-BF_3^-Cat^+$, $-Si(OR^{18})(OR^{19})(OR^{20})$ or alkyl, where alkyl denotes a halogenated or unsubstituted alkyl radical having 1 to 15 C atoms, in which, in addition, one or more CH_2 groups may each be replaced, independently of one another, by $-C\equiv C-$, $-CH=CH-$, $-O-$, $-CO-$, $-CO-O-$ or $-O-CO-$ in such a way that O atoms are not linked directly to one another;

20

Y^{12} and Y^{13} , independently of one another, denote H or alkyl, where alkyl denotes a halogenated or unsubstituted alkyl radical having 1 to 15 C atoms, in which, in addition, one or more CH_2 groups may each be replaced, independently of one another, by $-C\equiv C-$, $-CH=CH-$, $-O-$, $-CO-$, $-CO-O-$ or $-O-CO-$ in such a way that O atoms are not linked directly to one another;

25

L^1 , L^2 and L^3 , independently of one another, denote H or F;

30

R^{12} and R^{13} , independently of one another, denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms or toge-

ther form a $-(CH_2)_p-$ unit, where $p = 2, 3, 4, 5$ or 6 , where one, two or three of these CH_2 groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms;

5

R^{14} denotes OH, O-aryl, O-aralkyl, O-alkyl, Cl, Br, aryl, aralkyl or alkyl;

10

R^{15} denotes aryl, aralkyl or a halogenated or unsubstituted alkyl radical having 1 to 15 carbon atoms, where, in addition, one or more CH_2 groups in this alkyl radical may each be replaced, independently of one another, by $-C\equiv C-$, $-CH=CH-$, $-O-$, $-CO-$, $-CO-O-$ or $-O-CO-$ in such a way that O atoms are not linked directly to one another;

15

R^{16} and R^{17} denote H or an unbranched or branched alkyl radical having 1 to 15 carbon atoms or together form a $-(CH_2)_p-$ unit, where $p = 2, 3, 4, 5$ or 6 , where one, two or three of these CH_2 groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms;

20

R^{18} , R^{19} and R^{20} , independently of one another, denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms;

25

Cat^+ is an alkali metal cation or a quaternary ammonium cation;

and

30


X^- is a weakly coordinating anion;

with the proviso

that W denotes >CH- if $b+d \neq 0$;

that Y^{11} does not denote =O, =C(SR¹²)(SR¹³) or =CF₂ if Y^{11} is con-

5



nected to B or D =


that Y^{11} denotes -H, -I, -OH, -SH, -CO₂R¹⁴, -OSO₂R¹⁵,

10

-C(=S⁺R¹²)(SR¹³)X⁻, -B(OR¹⁶)(OR¹⁷), -BF₃⁻Cat⁺, -Si(OR¹⁸)(OR¹⁹)(OR²⁰)

or alkyl, where alkyl denotes a halogenated or unsubstituted alkyl radical having 1 to 15 C atoms, in which one or more CH₂ groups have each been replaced, independently of one another, by -C≡C-, -CH=CH-, -O-, -CO-, -CO-O- or -O-CO- in such a way that O atoms are not linked directly to one another and alkyl does not stand for


15



alkoxy, if W is connected directly to

where d is 0 or 1;

20




that B does not stand for if $d = 1$; and

that A can adopt identical or different meanings if a is 2.

25

2. Compound according to Claim 1, characterised in that

A stands for



30

3. Compound according to Claim 1, characterised in that

a is 0.

4. Compound according to any one of Claims 1 to 3, characterised in that
 Y^{12} and Y^{13} denote H.
- 5 5. Compound according to any one of Claims 1 to 4, characterised in that
 Z^{11} represents a single bond, $-CF_2O-$ or $-OCF_2-$.
- 10 6. Compound according to any one of Claims 1 to 5, characterised in that
 R^{11} denotes an unbranched halogenated or unsubstituted alkyl radical having 1 to 7 carbon atoms.
- 15 7. Compound according to any one of Claims 1 to 6, characterised in that
 Y^{11} denotes $=O$, $=C(SR^{12})(SR^{13})$ or $=CF_2$.
- 20 8. Compound according to any one of Claims 1 to 6, characterised in that
 Y^{11} denotes $-H$, $-F$, $-Cl$, $-Br$, $-I$, $-OH$, $-CO_2H$, $-C(=S^+R^{12})(-SR^{13})X^-$, $-B(OR^{16})(OR^{17})$, $-BF_3^-Cat^+$ or $-Si(OR^{18})(OR^{19})(OR^{20})$.
- 25 9. Compound according to any one of Claims 1 to 6 and 8, characterised in that
 X^- denotes BF_4^- , $CF_3SO_3^-$, $C_4F_9SO_3^-$, PF_6^- , SbF_6^- or AsF_6^- .
- 30 10. Compound according to any one of Claims 1 to 9, characterised in that
 b is 0 and d is 0.
11. Compound according to any one of Claims 1 to 9, characterised in that

characterised in that
a compound of the formula II




5

in which R^{11} , A, a and Z^{11} are as defined above for the formula IA,
is reacted in a reaction step (A1)

(A1) in the presence of a base with a compound of the formula III

10



15

in which Y^{12} and Y^{13} are as defined above for the formula IA, and R^{31}
denotes an alkyl radical having 1 to 15 carbon atoms, to give a com-
pound of the formula IV


20



25


in which R^{11} , A, a, Z^{11} , Y^{12} and Y^{13} are as defined above for the for-
mula IA, and R^{31} is as defined above for the formula III;
and subsequently, in a reaction step (A2),
(A2) the compound of the formula IV is converted into the com-
pound IA1

30



and optionally, in a reaction step (A3),


(A3) the compound of the formula IA1 is converted into the compound IA2



by reaction with CF_2Br_2 in the presence of $\text{P}(\text{N}(\text{R}^{21})_2)_3$,
 $\text{P}(\text{N}(\text{R}^{21})_2)_2(\text{OR}^{22})$ or $\text{P}(\text{N}(\text{R}^{21})_2)(\text{OR}^{22})_2$, where R^{21} and R^{22} , independently of one another, denote an alkyl radical having 1 to 15 carbon atoms;


or optionally, in a reaction step (A3'),

(A3') the compound of the formula IA1 is converted into the compound IA3




by reaction with $\text{CHG}(\text{SR}^{12})(\text{SR}^{13})$, in which G denotes $\text{P}(\text{OCH}_2\text{R}^{23})_3$, where R^{23} is a perfluorinated alkyl radical having 1 to 5 carbon atoms, or $\text{Si}(\text{CH}_3)_3$ or $\text{Si}(\text{CH}_2\text{CH}_3)_3$, and R^{12} and R^{13} are as defined above for the formula IA, in the presence of a strong base.

14. Process for the preparation of a compound of the formula IB



in which

R^{11} denotes H, F, Cl, Br, I, CN, aryl, heterocyclyl or alkyl;



a is 0, 1 or 2, where A can adopt identical or different meanings if a is 2;

Z^{11} represents a single bond, $-\text{CH}_2-\text{CH}_2-$, $-\text{CF}_2-\text{CF}_2-$, $-\text{CF}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CF}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-$, $-\text{CF}_2-\text{O}-$ or $-\text{O}-\text{CF}_2-$;

Y^{11} denotes $-\text{H}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CN}$, $-\text{OH}$ or $-\text{B}(\text{OR}^{16})(\text{OR}^{17})$;

Y^{12} and Y^{13} , independently of one another, denote H or alkyl;

L^1 , L^2 and L^3 , independently of one another, denote H or F; and

R^{16} and R^{17} , independently of one another, denote H or an un-

branched or branched alkyl radical having 1 to 15 carbon

atoms or together form a $-(\text{CH}_2)_p-$ unit, where $p = 2, 3, 4, 5$ or

6, where one, two or three of these CH_2 groups may be sub-


stituted by at least one unbranched or branched alkyl radical

having 1 to 8 carbon atoms;

characterised in that,


in a reaction step (B1),

(B1) a compound of the formula IA1




in which R^{11} , A, a, Z^{11} , Y^{12} and Y^{13} are as defined above for the formula IB,

is reacted with a compound of the formula V



V

in which L¹, L² and L³ are as defined above for the formula IB, M denotes Li, Cl-Mg, Br-Mg or I-Mg, and Q denotes H, F, Cl, Br, I or CN, with formation of the compound of the formula IB1




IB1

in which R¹¹, A, a, Z¹¹, Y¹², Y¹³, L¹, L² and L³ are as defined for the formula IB, and Q is as defined for the formula V;

and optionally, in a reaction step (B2),


(B2) the compound of the formula IB1 in which Q denotes Br is reacted with B(OR¹⁶)(OR¹⁷)(OR²⁴), where R¹⁶, R¹⁷ and R²⁴ are an unbranched or branched alkyl radical having 1 to 15 carbon atoms, or with HB(OR¹⁶)(OR¹⁷), where R¹⁶ and R¹⁷ denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms or together form a -(CH₂)_p- unit, where p = 2, 3, 4, 5 or 6, where one, two or three of these CH₂ groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms, in the presence of an alkyllithium base, to give the compound of the formula IB2



5

and optionally, in a reaction step (B3),

(B3) the compound IB2 is converted into the compound IB3




10

by reaction with an aqueous acid;

and/or optionally, in a reaction step (B4),

(B4) the compound IB2 or the compound IB3 is converted into the compound IB4


15



20

by reaction with hydrogen peroxide in alkaline or acidic solution.

15. Process for the preparation of a compound of the general formula IC




25

in which

R^{11} denotes H, F, Cl, Br, I, CN, aryl, heterocyclyl or alkyl;

30



a is 0, 1 or 2, where A can adopt identical or different meanings if a is 2;

Z^{11} represents a single bond, $-\text{CH}_2-\text{CH}_2-$, $-\text{CF}_2-\text{CF}_2-$, $-\text{CF}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CF}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-$, $-\text{CF}_2-\text{O}-$ or $-\text{O}-\text{CF}_2-$;


Y^{11} denotes $=\text{O}$, $=\text{C}(\text{SR}^{12})(\text{SR}^{13})$ or $=\text{CF}_2$;

Y^{12} and Y^{13} , independently of one another, denote H or alkyl; and


R^{12} and R^{13} , independently of one another, denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms or together form a $-(\text{CH}_2)_p-$ unit, where $p = 2, 3, 4, 5$ or 6 , where one, two or three of these CH_2 groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms;

characterised in that, in a reaction step (C1),

(C1) the compound of the formula IB4




in which R^{11} , A, a, Z^{11} , Y^{12} and Y^{13} are as defined above for the formula IC, and L^1 , L^2 and L^3 denote H, is converted into the compound IC1



5

using hydrogen in the presence of a transition-metal catalyst;
and optionally, in a reaction step (C2),

(C2) the compound IC1 is converted into the compound IC2




10

by reaction with CF_2Br_2 in the presence of $\text{P}(\text{N}(\text{R}^{21})_2)_3$,
 $\text{P}(\text{N}(\text{R}^{21})_2)_2(\text{OR}^{22})$ or $\text{P}(\text{N}(\text{R}^{21})_2)(\text{OR}^{22})_2$, where R^{21} and R^{22} , independently of one another, are an alkyl radical having 1 to 15 carbon atoms;
or optionally, in a reaction step (C2'),

15

(C2') the compound of the formula IC1 is converted into the compound IC3




20

by reaction with $\text{CHG}(\text{SR}^{12})(\text{SR}^{13})$, in which G denotes $\text{P}(\text{OCH}_2\text{R}^{23})_3$,
where R^{23} is a perfluorinated alkyl radical having 1 to 5 carbon atoms,
or $\text{Si}(\text{CH}_3)_3$ or $\text{Si}(\text{CH}_2\text{CH}_3)_3$, and R^{12} and R^{13} are as defined above for
the formula IC, in the presence of a strong base.

25

16. Process for the preparation of a compound of the formula ID




30

in which

R^{11} denotes H, F, Cl, Br, I, CN, aryl, heterocyclyl or alkyl;

5

A stands for



a is 0, 1 or 2, where A can adopt identical or different meanings if a is 2;

10

Z^{11} represents a single bond, $-\text{CH}_2-\text{CH}_2-$, $-\text{CF}_2-\text{CF}_2-$, $-\text{CF}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CF}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-$, $-\text{CF}_2-\text{O}-$ or $-\text{O}-\text{CF}_2-$;

Y^{11} denotes $-\text{CO}_2\text{H}$ or $-\text{C}(=\text{S}^+\text{R}^{12})(-\text{SR}^{13})\text{X}^-$;

Y^{12} and Y^{13} , independently of one another, denote H or alkyl;

15

L^1 , L^2 and L^3 , independently of one another, denote H or F;

R^{12} and R^{13} , independently of one another, denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms or together form a $-(\text{CH}_2)_p-$ unit, where $p = 2, 3, 4, 5$ or 6 , where one, two or three of these CH_2 groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms; and


20

X^- is a weakly coordinating anion;

characterised in that, in a reaction step (D1),

(D1) a compound of the formula IB1

25




IB1

30

in which R^{11} , A, a, Z^{11} , Y^{12} , Y^{13} , L^1 , L^2 and L^3 are as defined for the formula ID, and Q denotes H or Br,

is reacted with an organometallic base and CO_2 to give the compound ID1

5




10

in which R^{11} , A, a, Z^{11} , Y^{12} , Y^{13} , L^1 , L^2 and L^3 are as defined for the formula ID;

and optionally, in a reaction step (D2),

(D2) the compound ID1 is converted into the compound ID2

15




20

in the presence of an acid HX using HSR^{12} and HSR^{13} or using $HSR^{12}R^{13}SH$.

17. Process for the preparation of a compound of the formula IE


25



in which

R^{11} denotes H, F, Cl, Br, I, CN, aryl, heterocyclyl or alkyl;

30



a is 0, 1 or 2, where A can adopt identical or different meanings if a is 2;

Z^{11} represents a single bond, $-\text{CH}_2-\text{CH}_2-$, $-\text{CF}_2-\text{CF}_2-$, $-\text{CF}_2-\text{CH}_2-$, $-\text{CH}_2-\text{CF}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{O}-\text{CH}_2-$, $-\text{CF}_2-\text{O}-$ or $-\text{O}-\text{CF}_2-$;

Y^{11} denotes $-\text{CO}_2\text{H}$ or $-\text{C}(=\text{S}^+\text{R}^{12})(-\text{SR}^{13})\text{X}^-$;


Y^{12} and Y^{13} , independently of one another, denote H or alkyl;

R^{12} and R^{13} , independently of one another, denote an unbranched or branched alkyl radical having 1 to 15 carbon atoms or together form a $-(\text{CH}_2)_p-$ unit, where $p = 2, 3, 4, 5$ or 6 , where one, two or three of these CH_2 groups may be substituted by at least one unbranched or branched alkyl radical having 1 to 8 carbon atoms; and


X^- is a weakly coordinating anion;

characterised in that, in a reaction step (E1),

(E1) the compound of the formula ID1



in which R^{11} , A, a, Z^{11} , Y^{12} and Y^{13} are as defined above for the formula IE, and L^1 , L^2 and L^3 denote H, is converted into the compound IE1




5

using hydrogen in the presence of a transition-metal catalyst;
and optionally, in a reaction step (E2),

(E2) the compound of the formula IE1 is converted into the compound IE2

10



in the presence of an acid HX using HSR¹² and HSR¹³ or using HSR¹²R¹³SH.

15

20

25

30